

## **$\alpha$ 7 NICOTINIC RECEPTOR PRESENT ON MOUSE CORTICAL ASTROCYTES MODULATES ENDOGENEOUS GLUTAMATE RELEASE**

Patti Laura<sup>1</sup>, Raiteri Luca<sup>1</sup>, Grilli Massimo<sup>1</sup>, Robino Federica<sup>1</sup>, Zappettini Simona<sup>1</sup>,  
Bonanno Gian Battista<sup>1,2</sup>, Marchi Mario<sup>1,2</sup>

<sup>1</sup> Section of Pharmacology and Toxicology, Department of Experimental Medicine, University of Genoa, Italy

<sup>2</sup> Center of Excellence for Biomedical Research, University of Genoa, Italy

The presence of nicotinic receptors on astrocytes in human and rat brain has been previously demonstrated (1) however their possible functional role is still poorly understood. In this study we investigated on the presence of nicotinic receptors on gliosomes (2), purified from mouse cortex, and on their role in eliciting glutamate release. Epibatidine significantly increased basal release of [<sup>3</sup>H]D-aspartate and of endogenous glutamate from mouse gliosomes but not from synaptosomes. This effect was prevented by methyllycaconitine,  $\alpha$ -bungarotoxin and mecamylamine but not by dihydro-b-erythroidine. Epibatidine provoked also a significant increase of calcium concentration in gliosomes but not in synaptosomes; the [ $\text{Ca}^{2+}$ ]<sub>i</sub> enhancement induced by epibatidine and KCl in gliosomes were very similar each other. The present results indicate that  $\alpha$ 7 nicotinic receptors exist on mouse cortical glial particles and stimulate glutamate release.

(1) Xiu J, Nordberg A., Zhang JT., Guan ZZ., (2005) *Neurochem.Int.* 47, 281-290

(2) Stigliani S. et al (2006). *J Neurochem.* Feb, 96(3):656-68.

This work was supported by a MIUR Network grant.